



Rabies

(to be carried at all times and presented to medical personnel in the event of exposure)

FRONT

Rabies

IMPORTANT INFORMATION FOR EMERGENCY PERSONNEL

The individual carrying this card works in a laboratory with Rabies and may have come into contact via infectious droplets and aerosols containing rabies virus.

SPECIAL HAZARDS: Fixed tissue preparations can still be infectious, so extreme care is needed when handling them.

Note: This card is intended to be informational and should not substitute for clinical presentation and provider judgment of individual clinical cases.

BACK

MEDICAL ALERT INFORMATION

The person carrying this card has an occupational exposure to Rabies.
Immediately call Occupational Health at [301.405.3315](tel:301.405.3315) and consult with Infectious Diseases

COMMON SYMPTOMS

- Incubation period: days to years
- Initial presentation:
 - Flu-like symptoms, apprehension, behavior changes, headache, fever, malaise
 - Progressive encephalomyelitis, usually fatal

POST-EXPOSURE TREATMENT

- Thoroughly wash any skin wounds with soap and water and/or flush eyes or other affected mucous membranes with water
- Baseline serology: Rabies antibody
- Repeat rabies vaccination (even if employee has already been immunized)
 - If no past immunization, also use Rabies immune globulin (RIG)